

ON THE HOMOLOGICAL DIMENSION OF O-MINIMAL AND SUBANALYTIC SHEAVES

Abstract

Here we prove that the homological dimension of the category of sheaves on a topological space satisfying some suitable conditions is finite. In particular, we find conditions to bound the homological dimension of o-minimal and subanalytic sheaves.

1 Introduction

In [9] we studied the category $\text{Mod}(k_{X_{sa}})$ of sheaves on the subanalytic site X_{sa} associated to a real analytic manifold X . We defined the subcategory of quasi-injective sheaves (i.e. F is quasi-injective if the restriction $\Gamma(U; F) \rightarrow \Gamma(V; F)$ is surjective for each $U \supseteq V$ open subanalytic relatively compact) and we saw that quasi-injective are injective with respect to the functors of direct image, proper direct image and $\mathcal{H}om(G, \cdot)$ when G is a $\mathbb{R}$ -constructible sheaf on X . Moreover we proved that the quasi-injective dimension of $\text{Mod}(k_{X_{sa}})$ is finite, which implies that the cohomological dimension of the above functors is finite. However we had no answer concerning the homological dimension of $\text{Mod}(k_{X_{sa}})$ (see Remark 2.3.5 of [9]). The aim of this paper is to show that under some conditions (concerning the cardinality $2^{\aleph_0}$ of the reals) it is possible to bound this dimension. The key point is the fact that locally the category of subanalytic sheaves is equivalent to the category of sheaves on a topological space $\tilde{X}$. We are reduced to work with k -sheaves on a topological space, and in this case the homological dimension is equal to the flabby dimension. Hence we are reduced to bound the cohomological dimension of $\Gamma(U; \cdot)$, for any open subset U of $\tilde{X}$. In order to do that we need to assume that the cardinality of subanalytic subsets (which is equal to $2^{\aleph_0}$) is smaller than $\aleph_k$, $k < \infty$.

In more details the contents of this paper are as follows.

In § 2 we recall some notions as the definitions of injective and flabby sheaf and the homological and flabby dimension of the category of sheaves. We refer to [6] for more details.

In § 3 we study the general case of sheaves on a topological space X with a basis $\mathcal{T}$ whose elements are Lindelöf and stable under finite unions and intersections. We define the subcategory of $\mathcal{T}$ -flabby sheaves on X by saying that F is $\mathcal{T}$ -flabby if the restriction $\Gamma(U; F) \rightarrow \Gamma(V; F)$ is surjective for each $U, V \in \mathcal{T}$ with $U \supseteq V$. Since $\mathcal{T}$ forms a basis of the topology of X , we have that each open U of X has a covering $\{U_i\}_{i \in I}$ with $U_i \in \mathcal{T}$. Hence

$$R\Gamma(U; F) \simeq \varprojlim_{i \in I} R\Gamma(U_i; F),$$

and to bound the cohomological dimension of the functor $\Gamma(U; \cdot)$ is sufficient to bound the cohomological dimension of the functors $\Gamma(V; \cdot)$, $V \in \mathcal{T}$ and the cohomological dimension of the projective limit. The cohomological dimension of the projective limit is bounded if the cardinality of the index set I is smaller than $\aleph_k$ for some $k < \infty$, (see [8] or [10] in the more general setting of quasi-abelian categories). Hence it is bounded if the cardinality of $\mathcal{T}$ is smaller than $\aleph_k$, $k < \infty$.

In § 4 we consider an o-minimal structure $\mathcal{M} = (M, <, \dots)$. Let $\tilde{X}$ be the o-minimal spectrum of a definable space X . In this case $\mathcal{T}$ is the family of open constructible subsets of $\tilde{X}$. In the case of an o-minimal expansion of an ordered group, by a result of [4], the $\mathcal{T}$ -flabby dimension of $\tilde{X}$ is finite. Moreover the cardinality of $\mathcal{T}$ is bounded by the product of the cardinality of M and the cardinality of the language of the structure $\mathcal{M}$.

The case of subanalytic sheaves on a real analytic manifold is studied in § 5. We first reduce to $\text{Mod}(k_{U_{X_{sa}}})$ where U is a relatively compact subanalytic subset of X isomorphic to $\mathbb{R}^N$ endowed with the Grothendieck topology induced by X . This category is equivalent to $\text{Mod}(k_{\tilde{U}})$, where $\tilde{U}$ is the o-minimal spectrum of U . In this case $\mathcal{T}$ is the family of open globally subanalytic subsets of U , and its cardinality is $2^{\aleph_0}$. Hence if we assume that $2^{\aleph_0}$ is smaller or equal than $\aleph_k$, $k < \infty$ (in the case $k = 1$ this is the continuum hypothesis), then the homological dimension of $\text{Mod}(k_{\tilde{U}})$ (and hence of $\text{Mod}(k_{U_{X_{sa}}})$) is bounded.

We end this work with an example showing that in $\tilde{X}$ there are open subsets which do not admit countable covers.

2 Some preliminaries

We introduce some fundamental results about sheaves we will use in the rest of the paper. We refer to [6] for more details. Let X be a topological space and let k be a field. As usual, we denote by $\text{Mod}(k_X)$ the category of sheaves of k -vector spaces.

Definition 2.1 *Let $F \in \text{Mod}(k_X)$.*

- *F is injective if the functor $\text{Hom}(\cdot, F)$ is exact on $\text{Mod}(k_X)$.*
- *F is flabby if for any open subset U of X the restriction morphism $\Gamma(X; F) \rightarrow \Gamma(U; F)$ is surjective.*

In general injective $\Rightarrow$ flabby. When k is a field we have injective $\Leftrightarrow$ flabby ([6], Exercise II.10).

Definition 2.2 *The homological (resp. flabby) dimension of the category $\text{Mod}(k_X)$ is the smallest $N \in \mathbb{N} \cup \{\infty\}$ such that for any $F \in \text{Mod}(k_X)$ there exists an exact sequence*

$$0 \rightarrow F \rightarrow I^0 \rightarrow \dots \rightarrow I^N \rightarrow 0$$

with I^j injective (resp. flabby) for $0 \leq j \leq N$.

We shall need the following results.

- The homological dimension of $\text{Mod}(k_X)$ is finite if and only if there exists $N \in \mathbb{N}$ such that $R^j \text{Hom}(G, F) = 0$ for any $F, G \in \text{Mod}(k_X)$ and any $j > N$.
- The flabby dimension of $\text{Mod}(k_X)$ is finite if and only if there exists $N \in \mathbb{N}$ such that for any open subsets U of X , any $F \in \text{Mod}(k_X)$ and any $j > N$ we have $R^j \Gamma(U; F) = 0$ (i.e. if the functor $\Gamma(U; \cdot)$ has finite cohomological dimension).

In particular, when k is a field, since injective $\Leftrightarrow$ flabby, the homological dimension is finite if and only if the functor $\Gamma(U; \cdot)$ has finite cohomological dimension for any open subsets U of X .

3 $\mathcal{T}$ -flabby sheaves and homological dimension

Definition 3.1 A Lindelöf space is a topological space in which every open cover has a countable subcover.

Definition 3.2 The Lindelöf degree $l(X)$ of a topological space X is the smallest cardinal κ such that every open cover of the space X has a subcover of size at most κ . In this notation, X is Lindelöf iff $l(X) = \aleph_0$.

Let k be a field. Let X be a topological space and suppose that X admits a family $\mathcal{T}$ of open subsets such that

T1: each element of $\mathcal{T}$ is Lindelöf,

T2: $\mathcal{T}$ is stable under finite unions and intersections,

T3: $\mathcal{T}$ forms a basis for the topology of X .

Definition 3.3 A sheaf $F \in \text{Mod}(k_X)$ is said to be $\mathcal{T}$ -flabby if the restriction morphism $\Gamma(U; F) \rightarrow \Gamma(V; F)$ is surjective for each $U, V \in \mathcal{T}$ with $V \subseteq U$.

Remark that flabby $\Rightarrow \mathcal{T}$ -flabby.

Definition 3.4 The $\mathcal{T}$ -flabby dimension of $\text{Mod}(k_X)$ is the smallest $N \in \mathbb{N} \cup \infty$ such that for any $F \in \text{Mod}(k_X)$ there exists an exact sequence

$$0 \rightarrow F \rightarrow I^0 \rightarrow \cdots \rightarrow I^N \rightarrow 0$$

with I^j $\mathcal{T}$ -flabby for $0 \leq j \leq N$.

Proposition 3.5 Let $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ be an exact sequence in $\text{Mod}(k_X)$ with F' $\mathcal{T}$ -flabby. Then for any open subset U which is Lindelöf the morphism

$$\Gamma(U; F) \rightarrow \Gamma(U; F'')$$

is surjective.

Proof. We first consider a section $s'' \in \Gamma(U; F'')$. Since $F \rightarrow F''$ is surjective, we may find a covering $U = \bigcup_{i \in I} U_i$, $U_i \in \mathcal{T}$ and $s_i \in \Gamma(U_i; F)$ whose image is $s''|_{U_i}$. Since U is Lindelöf, we may find a countable subcover $U = \bigcup_{n \in \mathbb{N}} U_n$, $U_n \in \mathcal{T}$ and $s_n \in \Gamma(U_n; F)$ whose image is $s''|_{U_n}$.

Set $V_n = \bigcup_{i=1}^n U_i$. We prove by induction on n that there exists a section $t_{n+1} \in \Gamma(V_{n+1}; F)$ whose image is $s''|_{V_{n+1}}$ and $t_{n+1}|_{V_n} = t_n$.

This is clear for $n = 0, 1$ and we assume it is proved for n . By the induction hypothesis there exists a section $t_n \in \Gamma(V_n; F)$ whose image is $s''|_{V_n}$ and $t_n|_{V_m} = t_m$ if $m < n$. We set for short $V_n = W_1$ and $U_{n+1} = W_2$. We have seen that there exist $t_j \in \Gamma(W_j; F)$ whose image is $s''|_{W_j}$ for $j = 1, 2$. On $W_1 \cap W_2$ $t_1 - t_2$ defines a section of $\Gamma(W_1 \cap W_2; F')$ which extends to $t' \in \Gamma(V_{n+1}; F')$ because F' is $\mathcal{T}$ -flabby. Replace t_2 with $t_2 + t'$. We may suppose that $t_1 = t_2$ on $W_1 \cap W_2$. Then there exists $t_{n+1} \in \Gamma(V_{n+1}; F)$ such that $t_{n+1}|_{W_j} = t_j$, $j = 1, 2$. The t_n 's glue together into a section

$$s \in \varprojlim_{n \in \mathbb{N}} \Gamma(V_n; F) \simeq \Gamma(U; F)$$

which is sent to s'' , which proves the surjectivity of the morphism. $\square$

Proposition 3.6 *Let $F', F, F'' \in \text{Mod}(k_X)$, and consider the exact sequence*

$$0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0.$$

Suppose that F' is $\mathcal{T}$ -flabby. Then F is $\mathcal{T}$ -flabby if and only if F'' is $\mathcal{T}$ -flabby.

Proof. Let $U, V \in \mathcal{T}$ with $V \subseteq U$ and let us consider the diagram below

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \Gamma(U; F') & \longrightarrow & \Gamma(U; F) & \longrightarrow & \Gamma(U; F'') & \longrightarrow & 0 \\ & & \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \\ 0 & \longrightarrow & \Gamma(V; F') & \longrightarrow & \Gamma(V; F) & \longrightarrow & \Gamma(V; F'') & \longrightarrow & 0 \end{array}$$

where the row are exact by Proposition 3.5 and the morphism α is surjective since F' is $\mathcal{T}$ -flabby. It follows from the five lemma that β is surjective if and only if γ is surjective. $\square$

Proposition 3.7 *Let F be $\mathcal{T}$ -flabby. Then F is $\Gamma(U; \cdot)$ -injective for each open $U \subseteq X$ which is Lindelöf.*

Proof. The family of $\mathcal{T}$ -flabby sheaves contains injective sheaves, hence it is cogenerating. Then the result follows from Propositions 3.5 and 3.6. $\square$

Now let us consider the case of open subsets of Lindelöf degree $\aleph_k$, $k < \infty$. In order to do that we need the following result.

Proposition 3.8 *Let X be a topological space admitting a family $\mathcal{T}$ satisfying T1-T3. Suppose that X has finite $\mathcal{T}$ -flabby dimension. Let U be an open subset of X of Lindelöf degree $\aleph_k$, $k < \infty$. Then the cohomological dimension of $\Gamma(U; \cdot)$ is finite.*

Proof. We have

$$R\Gamma(U; F) \simeq R\varprojlim_{i \in I} R\Gamma(U_i; F)$$

where $U = \bigcup_{i \in I} U_i$ with $U_i \in \mathcal{T}$ and $\sharp I$ smaller than $\aleph_k$. Since X has finite $\mathcal{T}$ -flabby dimension and using Proposition 3.7 for each i we may replace $R\Gamma(U_i; F)$ with $\Gamma(U_i; I^\bullet)$, where $I^\bullet$ is a $\mathcal{T}$ -flabby resolution of F of length $N < \infty$. Since cohomological dimension of $\varprojlim_{i \in I}$ is finite if $\sharp I$ is smaller than $\aleph_k$, $k < \infty$ (see [8, 10]), then the j -th cohomology of $R\varprojlim_{i \in I} R\Gamma(U_i; F)$ vanishes for $j > N + M$. Since $M, N < \infty$ are independent of F and i , the result follows. $\square$

Theorem 3.9 *Let X be a topological space admitting a family $\mathcal{T}$ satisfying T1-T3. Suppose that X has finite $\mathcal{T}$ -flabby dimension. Suppose that there exists $k < \infty$ such that every open subset of X has Lindelöf degree $\aleph_k$. Then $\text{Mod}(k_X)$ has finite homological dimension.*

Proof. By Proposition 3.8 X has finite flabby dimension. In the case of sheaves of k -vector spaces we have flabby $\Leftrightarrow$ injective and the result follows. $\square$

4 Homological dimension of o-minimal sheaves

Let $\mathcal{M} = (M, <, \dots)$ be an o-minimal structure. In [4] the authors studied sheaf cohomology of sheaves on $\tilde{X}$, the o-minimal spectrum of a definable space X . The category of sheaves on $\tilde{X}$ is equivalent to the category of sheaves on the o-minimal site, consisting of open definable subsets of X and coverings admitting a finite refinement.

Let $\mathcal{T}$ be the family of open constructible subsets of $\tilde{X}$. Then $\mathcal{T}$ satisfies T1-T3, indeed

- $\mathcal{T}$ forms a basis for the topology of $\tilde{X}$,

- every element of $\mathcal{T}$ is quasi-compact,
- $\mathcal{T}$ is stable under finite unions and intersections.

For each open subset U of $\tilde{X}$ and each $F \in \text{Mod}(k_{\tilde{X}})$ we have

$$R\Gamma(U; F) \simeq \varprojlim_{i \in I} R\Gamma(U_i; F)$$

where $U = \bigcup_{i \in I} U_i$ and $U_i \in \mathcal{T}$ for each $i \in I$. In order to apply Theorem 3.9 we need that:

- the $\mathcal{T}$ -flabby dimension of $\tilde{X}$ is finite,
- $\sharp I \leq \aleph_k, k < \infty$.

If every open constructible subset U of X is normal (this is true in the case of an o-minimal expansion of an ordered group), then by Proposition 4.2 of [4] we have $R^j\Gamma(U; F) = 0$ for all $j > \dim U = \dim X$.

Concerning $\sharp I$, we have that $\sharp I \leq \sharp \mathcal{T}$, and $\sharp \mathcal{T} \leq \sharp \mathcal{L} \cup M = \sharp \mathcal{L} \cdot \sharp M$, where $\mathcal{L}$ denotes the language of the structure $\mathcal{M}$.

5 Homological dimension of subanalytic sheaves

Let X be a real analytic manifold, let X_{sa} the associated subanalytic site and denote by $\text{Mod}(k_{X_{sa}})$ the category of sheaves of k -vector spaces on X_{sa} . We refer to [7, 9] for the theory of subanalytic sheaves. We will see that under suitable hypothesis the homological dimension of $\text{Mod}(k_{X_{sa}})$ is finite, i.e. there exists $N \in \mathbb{N}$ such that $R^j\text{Hom}(G, F) = 0$ for any $F, G \in \text{Mod}(k_{X_{sa}})$ and any $j > N$.

Let X be a real analytic manifold. Then X has an atlas $(U_n, \phi_n)_{n \in \mathbb{N}}$, $\phi_n : U_n \xrightarrow{\sim} \mathbb{R}^M$, U_n relatively compact open subanalytic subset. Let us consider the subanalytic site X_{sa} associated to X . In order to bound $\text{RHom}(F, G)$ for any $F, G \in \text{Mod}(k_{X_{sa}})$ it suffices to bound $R\mathcal{H}om(F, G)$. Indeed we have $\text{RHom}(F, G) \simeq R\Gamma(X; R\mathcal{H}om(F, G))$ and $\Gamma(X; \cdot)$ has finite cohomological dimension (see [9]).

We have $R^j\mathcal{H}om(F, G) = 0$ if $R^j\mathcal{H}om(F, G)|_{U_n} = 0$ for each $n \in \mathbb{N}$. Then we may reduce to the case U_n with the Grothendieck topology induced by the subanalytic site, i.e. the category of globally subanalytic open subsets of $\mathbb{R}^M$ with coverings admitting a finite subcover. Globally subanalytic geometry is defined by the o-minimal structure given by the ordered field of real numbers expanded by globally analytic functions $\mathbb{R}_{\text{an}} = (\mathbb{R}, <, 0, 1, +, \cdot, (f)_{f \in \text{an}})$,

where f is a restriction to $[-1, 1]^m$ of a convergent power series on some neighborhood of $[-1, 1]^m$.

Set $X = U_n$ and let $\tilde{X}$ be the o-minimal spectrum of X . The family $\mathcal{T}$ of open sets $\tilde{U}$ where U is open globally subanalytic satisfies T1-T3. Moreover the $\mathcal{T}$ -flabby dimension of $\text{Mod}(k_{\tilde{X}})$ is finite. Indeed by Proposition 4.2 of [4] the cohomological dimension of $\Gamma(\tilde{U}; \cdot)$ is finite for each $\tilde{U} \in \mathcal{T}$.

Let us see that $\sharp \mathcal{T} \leq 2^{\aleph_0}$. In order to see that we have to show that $\sharp \mathcal{L} = 2^{\aleph_0}$, where $\mathcal{L}$ denotes the language of $\mathbb{R}_{\text{an}}$. This is true if $\sharp \{(f)_{f \in \text{an}}\} = 2^{\aleph_0}$. This is seen by identifying $f = \sum_{I \in \mathbb{N}^m} a_I x^I$ with $(a_I)_{I \in \mathbb{N}^m} \in \mathbb{R}^{\mathbb{N}^m}$. Then $\sharp \{(f)_{f \in \text{an}}\} \leq \sharp \mathbb{R}^{\mathbb{N}^m} = (2^{\aleph_0})^{\aleph_0} = 2^{\aleph_0 \cdot \aleph_0} = 2^{\aleph_0}$ (see [5]).

Let U be an open subset of $\tilde{X}$. Then $U = \bigcup_{i \in I} U_i$ with $U_i \in \mathcal{T}$ and $\sharp I$ is smaller than $\sharp \mathcal{T} = 2^{\aleph_0}$. Hence every open subset of X has Lindelöf degree at most $2^{\aleph_0}$. If we assume that $2^{\aleph_0}$ is smaller than $\aleph_k$ for some $k < \infty$ (if we suppose $\leq \aleph_1$ it is the continuum hypothesis), then Theorem 3.9 implies that the homological dimension of $\text{Mod}(k_{\tilde{X}})$ is finite.

Remark 5.1 *In $\tilde{X}$ there are open subsets which are not Lindelöf. For example let us consider the open set (in $\tilde{X} = \widetilde{\mathbb{R}^2}$ with respect to $\mathbb{R}_{\text{an}}$)*

$$U = \bigcup_{r \in (0,1) \setminus \mathbb{Q}} \tilde{V}_r,$$

where

$$V_r = \{(x, y) \in \mathbb{R}^2 ; 0 < x < 1, r < y < r + x\}.$$

We prove that $\{V_r\}_{r \in (0,1) \setminus \mathbb{Q}}$ has no countable subcover.

We argue by contradiction. Suppose that there exists a countable subcover $\{\tilde{V}_{r_n}\}_{n \in \mathbb{N}}$ of U in $\tilde{X}$. Let 0^+ be the ultrafilter defined by $0^+ = \{S \supset (0, \varepsilon)\}$, where $\varepsilon > 0$ and S is globally subanalytic. Let $\pi : \mathbb{R}^2 \rightarrow \mathbb{R}$ be the projection onto the first coordinate. We have

$$U \cap \tilde{\pi}^{-1}(0^+) = \bigsqcup_{r \in (0,1) \setminus \mathbb{Q}} (\tilde{V}_r \cap \tilde{\pi}^{-1}(0^+)).$$

Indeed, let $x \in \tilde{V}_r \cap \tilde{\pi}^{-1}(0^+)$, and let $s \neq r$. Let $\varepsilon < |r - s|$, then $V_s \cap \pi^{-1}((0, \varepsilon)) \cap V_r = \emptyset$ and $\tilde{V}_s \cap \tilde{\pi}^{-1}(0^+) \subset \tilde{V}_s \cap \tilde{\pi}^{-1}(\widetilde{(0, \varepsilon)})$ since $0^+ \subset \widetilde{(0, \varepsilon)}$ for any $\varepsilon > 0$. Hence $\tilde{V}_r \cap \tilde{\pi}^{-1}(0^+) \cap \tilde{V}_s = \emptyset$ if $r \neq s$. Moreover

$$\bigsqcup_{r \in (0,1) \setminus \mathbb{Q}} (\tilde{V}_r \cap \tilde{\pi}^{-1}(0^+)) \subset \bigcup_{n \in \mathbb{N}} \tilde{V}_{r_n} = U.$$

Since $\sharp((0, 1) \setminus \mathbb{Q}) = 2^{\aleph_0} > \aleph_0$, there exists $n \in \mathbb{N}$ such that

$$\sharp \left\{ t \in (0, 1) \setminus \mathbb{Q} ; \tilde{V}_t \cap \tilde{\pi}^{-1}(0^+) \subseteq \tilde{V}_{r_n} \right\} \geq 2.$$

But $\tilde{V}_t \cap \tilde{\pi}^{-1}(0^+) \subseteq \tilde{V}_{r_n}$ implies that $V_t = V_{r_n}$, so $t = r_n$ which is a contradiction.

Remark 5.2 More generally, as in Remark 5.1, given an *o-minimal expansion of an ordered group* $(\mathbb{R}, <, +, \dots)$, for any definable manifold X of dimension ≥ 2 , the space $\tilde{X}$ has open subsets which have Lindelöf degree $2^{\aleph_0}$.

Remark 5.3 It seems that the right setting is that of locally definable spaces in an *o-minimal structure*. In that setting the *o-minimal (definable)* case and the *subanalytic* case would be treated uniformly. However, to work on that setting one has to take into account the (special case of the) theory locally semi-algebraic spaces developed in [2, 3].

References

- [1] M. Coste *An introduction to o-minimal geometry* Dip. Mat. Univ. Pisa, Dottorato di Ricerca in Matematica, Istituti Editoriali e Poligrafici Internazionali, Pisa (2000).
- [2] H. Delfs *Homology of locally semialgebraic spaces* Lecture Notes in Math. **1484** Springer-Verlag (1991).
- [3] H. Delfs and M. Knebusch *Locally semi-algebraic spaces* Lecture Notes in Math. **1173** Springer-Verlag (1985).
- [4] M. Edmundo, G. Jones and N. Peatfield *Sheaf cohomology in o-minimal structures* J. Math. Logic **6** pp. 163-179 (2006).
- [5] T. Jech *Introduction to set theory* Dekker (1984).
- [6] M. Kashiwara and P. Shapira *Sheaves on manifolds* Grundlehren der Math. **292** Springer Verlag (1990).
- [7] M. Kashiwara and P. Shapira *Ind-sheaves* Astérisque **271** (2001).
- [8] C. U. Jensen *Les foncteurs dérivés de $\varprojlim$ et leurs applications en théorie des modules* Lecture Notes in Math. **254** Springer-Verlag (1972).

- [9] L. Prelli *Sheaves on subanalytic sites* Rend. Sem. Mat. Univ. Padova **120** pp. 167-216 (2008).
- [10] F. Prosmans *Derived limits in quasi-abelian categories* Bull. Soc. Roy. Sci. Liège **68** pp. 335-401 (1999).